

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 072 253 A1**

(12) **EUROPEAN PATENT APPLICATION**
published in accordance with Art. 158(3) EPC

(43) Date of publication:
31.01.2001 Bulletin 2001/05

(21) Application number: 99913704.5

(22) Date of filing: 19.04.1999

(51) Int. Cl.⁷: **A61K 7/16**

(86) International application number:
PCT/JP99/02060

(87) International publication number:
WO 99/55296 (04.11.1999 Gazette 1999/44)

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

(30) Priority: 24.04.1998 JP 13141398
28.12.1998 JP 37387398
15.04.1999 JP 10780999

(71) Applicant:
SUNSTAR KABUSHIKI KAISHA
Takatsuki-shi, Osaka 569-1134 (JP)

(72) Inventors:
• TAKATSUKA, Tsutomu
Mishima-gun, Osaka 618-0001 (JP)
• NAKAO, Akira
Takatsuki-shi, Osaka 569-1142 (JP)

(74) Representative:
Stoner, Gerard Patrick et al
MEWBURN ELLIS
York House
23 Kingsway
London WC2B 6HP (GB)

(54) **ORAL COMPOSITIONS**

(57) Disclosed is an oral composition comprising palatinit. More particularly, disclosed is an oral composition comprising palatinit which exerts a synergistic effect when combined with a fluorine or zinc compound.

EP 1 072 253 A1

DescriptionTechnical Field

5 **[0001]** The present invention relates to an oral composition comprising palatinit, and more particularly, it relates to an oral composition comprising palatinit which exhibits the synergistic effect when combined with a fluorine or zinc compound.

Background Art

10

[0002] Dental caries is a state of a dental carious cavity caused by the dissolution of calcium from teeth and which can not naturally return to a healthy state. But, there is a state referred to as a sub-surface lesion which can return to the healthy state, in the course of the development of the dental carious cavity ("Zusetsu Ushoku-gaku" edited by Shoichi, Suga, 1990, 139). Therefore, in order to decrease dental caries, it is desirable to enhance the remineralization so that the teeth can return to the healthy state within a term during which the teeth return to the original state.

15

[0003] As a method for enhancing the remineralization, the use of fluorine compounds has been known for a long time. However, because the ingestion of a large amount of fluorine exhibits the toxicity, it is desired that fluorine be effectively utilized in an as smaller as possible amount. To accomplish this, the use of a substance that enhances the remineralization effect of fluorine is exemplified. For example, there is proposed a combination of fluorine and hydroxyapatite in JP-A-1-110608. However, its effect is insufficient and is not satisfactory.

20

[0004] On the other hand, a zinc compound that has been utilized in the oral composition generally has an astringent or salty taste and, therefore, there is a problem that it has the very bad feeling for use upon its use.

[0005] One object of the present invention is to provide an oral composition which has the high safety and can enhance the remineralization. Another object of the present invention is to provide an oral composition which exhibits the sufficient remineralization effect and has the better feeling for use.

25

Disclosure of Invention

30

[0006] In view of the above circumstances, the present inventors studied intensively the substances that can enhance the remineralization and, as a result, found that palatinit exhibits the excellent properties, that an oral composition comprising palatinit can enhance the remineralization, and, further, that the inclusion of palatinit in a combination with a fluorine or zinc compound can enhance the remineralization due to the synergistic effect of both ingredients, which resulted in the completion of the present invention.

35

[0007] Palatinit has hitherto been utilized in foods and the like as a low cariogenic sweetener. However, palatinit has never been utilized for an oral-use, and its remineralization effect is not known.

[0008] That is, the present invention has completed based on such novel findings, and, in the first aspect, provides an oral composition comprising palatinit which has the high safety and can enhance the remineralization.

[0009] Moreover, in the second aspect, the present invention provides an oral composition comprising palatinit and a remineralization enhancing ingredient, which exhibits the sufficient remineralization effect and has the better feeling for use.

40

[0010] The present invention is described below in detail according to the first and second aspects thereof in sequence.

[First aspect]

45

[0011] Palatinit to be used in the first and second aspects of the present invention is a sugar alcohol of a disaccharide, and may be α -D-glucopyranosyl-1,6-mannitol, its isomer, α -D-glucopyranosyl-1,6-sorbitol or a mixture thereof. Palatinit can be obtained by hydrogenation of palatinose which is converted from sucrose as a raw material with glycosyltransferase. And, palatinit is also a trade name of the product of Mitsui Sugar Co. Ltd. or Sudzucker A.G., and is also referred to as reduced-palatinose or isomalt. Palatinit is widely known as a non-cariogenic sugar which scarcely develops dental caries, based on the fact that the cariogenic microorganisms do not produce acids from palatinit in an oral cavity. Palatinit has been blended in non-sugar foods or a specified health food such as so-called "dental caries-resistant candy".

50

[0012] The amount of palatinit to be blended in the present oral composition is in the range of 0.1% to 60% by weight, preferably 1% to 40% by weight, based on the total weight of the oral composition. When the amount is less than 0.1% by weight, the desired effect can not be obtained. On the other hand, when the amount is more than 60% by weight, the stability of the formulation is deteriorated.

55

[Second aspect]

[0013] The remineralization enhancing ingredient to be used in the second aspect of the present invention is an ingredient which can remineralize the teeth from its sub-surface lesion state. Examples of the remineralization enhancing ingredient are fluorine compounds, zinc compounds, phosphorus compounds, calcium compounds and the like, but not limited thereto.

[0014] Examples of the fluorine compound to be used in the second aspect of the present invention as the remineralization enhancing ingredient are sodium fluoride, potassium fluoride, ammonium fluoride, stannous fluoride, sodium or potassium monofluorophosphate and the like. Particularly preferred are sodium fluoride and sodium monofluorophosphate.

[0015] These fluorine compounds, alone or in a combination thereof, may be blended in the present oral composition in the range of 0.1-5,000ppm, preferably 100-1,100ppm in terms of fluoride ion, based on the total weight of the present oral composition.

[0016] Further, the zinc compound to be used in the second aspect of the present invention as the remineralization enhancing ingredient is preferably a water-slightly soluble zinc compound, wherein the water-slightly soluble zinc compound is defined as having the solubility of less than 0.5g per 100g of water at 25°C, and water-insoluble zinc compound is included therein. Among them, particularly preferred are zinc oxide, zinc citrate and zinc stearate. From a viewpoint of a taste, the water-slightly soluble zinc compound having a smaller particle diameter and a greater specific surface area is preferred. More particularly, preferred are those having the particle diameter of not greater than 0.3 µm and the specific surface area of greater than 10m²/g. When the particle diameter exceeds 0.3 µm, the astringency becomes strong. Examples of the commercial products of the zinc compound are fine particle zinc white and hyperfine particle zinc oxide, "FINEX series" manufactured by Sakai Chemical Industry Co., Ltd. These water-slightly soluble zinc compounds, alone or in a combination thereof, may be blended in the oral composition in the amount of 0.1% to 5% by weight, based on the total weight of the present oral composition.

[0017] Further, when the oral composition of the second aspect of the present invention contains a zinc compound, preferred pH thereof is within the range of 6.0 to 8.5. When pH is lower than 6.0, then the astringency is strong, and when pH is higher than 8.5, then oral mucosa may be irritated, therefore, the composition out of the above pH range is not preferable.

[0018] Further, examples of the phosphorus compound to be used in the second aspect of the present invention as the remineralization enhancing ingredient are disodium hydrogenphosphate, sodium dihydrogenphosphate, dipotassium hydrogenphosphate, potassium dihydrogenphosphate, trisodium phosphate, tripotassium phosphate and the like, but not limited thereto.

[0019] Moreover, examples of the calcium compound to be used in the second aspect of the present invention as the remineralization enhancing ingredient are, for example, calcium chloride, calcium nitrate, calcium sulfate, calcium carbonate, calcium citrate, calcium hydrogenpyrophosphate, calcium gluconate, calcium glycerophosphate, calcium hydroxide, calcium oxide, calcium silicate and the like, but not limited thereto.

[0020] That is, the present invention provides an oral composition comprising palatinit alone or in a combination with any remineralization enhancing ingredients, which can enhance the remineralization due to the synergistic effect of both ingredients.

[0021] The oral composition of the present invention may be properly formulated, depending upon its use, into a form such as a toothpaste, powder or liquid dentifrice, wetting dentifrice, gel, cream, pasta, mouthwash, spray, foam, coating agent and the like, according to the conventional procedure. Other ingredients to be blended therein are not particularly limited, and the known active ingredients, polishing agents, humectants, thickening agents, foaming agents, preservatives, flavoring agents, sweeteners, pH adjusting agents, organic acids, sugar alcohol, anti-oxidizing agents and others known as the ingredient for the oral composition may be blended in the oral composition, so long as they do not deteriorate the effects of the present invention.

[0022] Examples of the active ingredient are enzymes such as amylase, protease, lysozyme and dextranase, the antimicrobial agents such as sanguinarine, allantoin, aminobenzoate derivatives, hexetidine, chlorhexidine, triclosan and cetylpyridinium chloride, vitamins such as vitamin B, C and E, and the astringents such as potassium nitrate, lithium nitrate and sodium nitrate.

[0023] Examples of the polishing agent are silica, alumina, aluminosilicate, aluminium hydroxide and the like.

[0024] Examples of the humectant are glycerol, propylene glycol, sorbitol, polyethylene glycol, polypropylene glycol and the like.

[0025] Examples of the thickening agent are sodium carboxymethyl cellulose, methyl cellulose, hydroxyethyl cellulose, alginates, xanthan gum, carrageenan, gum arabic, polyvinyl alcohol and the like.

[0026] Examples of the forming agent are anionic-, nonionic-, cationic- and amphoteric-surfactants. Examples of the anionic-surfactant are alkyl sulfates, sodium dodecylbenzene sulfonate, amino acids, sulfosuccinates, sucrose fatty acid esters and the like. Examples of the nonionic-surfactant are Pluronic series that are polyoxyethylene-polyoxypro-

pylene copolymer, fatty acid dialkanolamides and the like.

[0027] Examples of the preservative are methylparaben, propylparaben, benzoates, sodium benzoate, paraoxybenzoic acid esters, titanium dioxide and the like.

[0028] Examples of the flavoring agent are peppermint oil, spearmint oil, Japanese peppermint oil, orange oil, menthol, cloves oil, anise oil, wintergreen oil, eucalyptus oil and the like.

[0029] Examples of the sweetening agent or sweetener are saccharin salts, dextrose, Aspartame, xylitol, stevia extract, Acesulfame, granulated sugar, powdered sugar, starch syrup and the like. Although palatinit exhibits sweetness, the above sweetening agents or sweeteners may be added depending upon the feeling for use of the formulation.

[0030] Examples of the pH adjusting agent are citric acid and salts thereof, phosphoric acid and salts thereof; malic acid and salts thereof, gluconic acid and salts thereof, maleic acid and salts thereof; aspartic acid and salts thereof, gluconic acid and salts thereof; succinic acid and salts thereof, glucuronic acid and salts thereof, fumaric acid and salts thereof, glutamic acid and salts thereof, adipic acid and salts thereof, lactic acid and salts thereof, pantothenic acid and salts thereof, hydrochloric acid, alkali metal hydroxides and the like.

Brief Description of Drawings

[0031]

Figure 1 is a graph which compares the remineralization effects of the various sugars in the system not containing fluoride.

Figure 2 is a graph which compares the remineralization effects of the various sugars in the system containing fluoride.

The Best Mode for Carrying Out the Invention

[0032] The present invention will be further illustrated by way of the following Examples, which are not to be construed to limit the scope of the present invention. The amounts indicated in the Examples are all percents (%) by weight.

Experimental Example 1

Evaluation of the remineralizing ability 1

[0033] The remineralization effect of the sugars was evaluated in an *in vitro* test using a bovine tooth according to the procedures described in D. J. White *et al.*, *Caries Res.*, 21, 228 (1987).

1. An enamel section having 4mm length × 3mm width was obtained from the bovine tooth. The section was embedded in a dental resin to obtain an enamel block.
2. An enamel varnish was applied to a portion of approximately 1/3 of the surface of the enamel block in order not to cause the demineralization of that portion. Then, the enamel block was demineralized with a demineralizing solution containing 50% of saturated hydroxyapatite/0.1M lactic acid/pH 5.0 to prepare an artificial caries.
3. An enamel varnish was applied to a portion of approximately 2/3 of the surface of the treated enamel block in order not to cause the remineralization of that portion. The enamel block was immersed for ten days in a test solution prepared by adding the test sugar into an aqueous solution containing 3.0mM calcium ion and 1.8mM phosphate ion so that the concentration of the sugar became 20%, to perform the remineralization treatment.
4. Three thin sections having approximately 500 μm in thickness were prepared from the enamel block. The central portion of the thin sections was ground so that the thickness became approximately 100 μm using a double-side polishing machine.
5. The X-ray photograph was taken of the thin section prepared in the fourth step. The mineral amounts of the demineralized portion and the remineralized portion were calculated based on the brightness of the each portion and a distance from the surface using an image processor. The mineral amount difference between the demineralized and remineralized portions was expressed as a remineralization value. A greater value thereof shows the greater remineralization. A control test was performed in a similar manner to that described above, except that the test solution containing only 3.0mM calcium ion and 1.8mM phosphate ion but not containing the sugar was used.

[0034] The results are shown in Table 1 and Figure 1.

Table 1

Systems not containing fluoride	
Sugar (20%)	Remineralization value
Control	405
Sorbitol	645
Mannitol	640
Xylitol	416
Erythritol	602
Trehalose	715
Palatinit	1057
Unit: Vol. % μm	

[0035] Further, a similar test was performed in which sodium fluoride was added to the test solution to 2ppm in terms of fluoride ion.

[0036] The results are shown in Table 2 and Figure 2.

Table 2

Systems containing fluoride (2ppm)	
Sugar (20%)	Remineralization value
Control	1626
Sorbitol	1358
Mannitol	1377
Xylitol	1636
Erythritol	1330
Trehalose	1205
Palatinit	2374
Unit: Vol. % μm	

[0037] From Tables 1 and 2, it was found that, among the sugars tested, palatinit has the particularly excellent remineralizing ability and that the remineralizing ability is enhanced by combining palatinit with sodium fluoride.

Experimental Example 2

Evaluation of the remineralizing ability 2

[0038] The remineralization enhancing effects of the oral compositions to which the various sugars had been added were tested *in vitro* according to the similar manner to that described in Experimental Example 1.

[0039] Provided that, in the remineralization treatment, a solution containing 3.0mM calcium ion and 1.8mM phosphate ion to which a test dentifrice (see Table 3) had been added so as to form a 4-fold slurry was used, and the immersion period was set to be fourteen days. The results of the X-ray photograph were evaluated using the image processor according to a similar manner to that described in Experimental Example 1. The results are shown in Table 3.

Table 3

Ingredients	Example 1	Example 2	Example 3	Example 4	Example 5	Example 6	Comparative Example 1	Comparative Example 2	Comparative Example 3	Comparative Example 4	Comparative Example 5
Silicic acid anhydride	20	20	20	20	20	20	20	20	20	20	20
Sodium carboxymethyl cellulose	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5
Sorbitol	40	40	35	25	15	15	45	45	35	35	35
Sodium lauryl sulfate	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5
Saccharin sodium	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Sodium fluoride	-	0.2	0.2	0.2	0.2	0.2	-	0.2	0.2	0.2	0.2
Flavor	1	1	1	1	1	1	1	1	1	1	1
Xylitol	-	-	-	-	-	-	-	-	10	-	-
Erythritol	-	-	-	-	-	-	-	-	-	10	-
Trehalose	-	-	-	-	-	-	-	-	-	-	10
Palatinit	10	5	10	20	30	60	-	-	-	-	-
Remineralization value	852	1378	1468	1496	1548	1580	320	1024	958	842	862

From the above Table 3, it was found that palatinit has the particularly excellent remineralizing ability, also in the evaluation of Examples 1-6 and Comparative Examples 1-5.

Experimental Example 3

Evaluation of the remineralizing ability 3

5 [0040] The remineralization enhancing effect of the oral composition to which the zinc compound had been added was tested *in vitro* according to almost the same manner as that described in Experimental Example 1.

[0041] Provided that, in the remineralization treatment, a solution containing 3.0mM calcium ion and 1.8mM phosphate ion to which the test dentifrice (see Table 4) had been added so as to form a 4-fold slurry was used, and the pH cycling procedure was taken in which the enamel block was additionally immersed in the demineralization solution for
10 three hours per day. An evaluation period was set to be fourteen days. The results of the X-ray photograph were evaluated using the image processor according to a similar manner to that described in Experimental Example 1. The results are shown in Table 4.

15

20

25

30

35

40

45

50

55

Table 4

Ingredients	Example 7	Example 8	Example 9	Example 10	Example 11	Example 12	Example 13	Comparative Example 6	Comparative Example 7
Silicic acid anhydride	20	20	20	20	20	20	20	20	20
Sodium carboxymethyl cellulose	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5
Sorbitol	40	40	35	25	35	35	35	50	35
Sodium lauryl sulfate	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5
Saccharin sodium	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Sodium fluoride	-	0.2	0.2	0.2	0.2	0.2	0.2	-	0.2
Flavor	1	1	1	1	1	1	1	1	1
Xylitol	-	-	-	-	-	-	-	-	10
Palatinit	10	10	10	20	10	10	10	-	-
Zinc oxide	1	-	1	-	-	0.1	5	1	-
Zinc stearate	-	-	-	1	-	-	-	-	-
Zinc citrate	-	-	-	-	1	-	-	-	-
Remineralization value	102	513	542	516	630	530	721	10	354

[0042] From the above Table 4, in the evaluation of Examples 7-13 and Comparative Examples 6-7, it was shown that the remineralization is enhanced by blending palatinit and a zinc compound.

Experimental Example 4

Organoleptic evaluation of the astringency 1

- 5 [0043] The oral compositions of Examples 14-16 and Comparative Examples 8-9 in Table 5 were prepared. Ten healthy persons used these oral compositions according to the conventional manner, and the astringency after spout was organoleptically evaluated. The total points of ten healthy persons are shown according to the following 3 grades-criteria: 2: excellent; 1: good; 0: astringent.

[0044] The results are shown in Figure 5.

Table 5

Ingredients	Example 14	Example 15	Example 16	Comparative Example 8	Comparative Example 9
15 Silicic acid anhydride	20	20	20	20	20
20 Sodium carboxymethyl cellulose	1.5	1.5	1.5	1.5	1.5
Sorbitol	35	35	35	35	35
Sodium lauryl sulfate	1.5	1.5	1.5	1.5	1.5
25 Saccharin sodium	0.1	0.1	0.1	0.1	0.1
Sodium fluoride	0.2*	0.2	0.2	0.2	0.2
Flavor	1	1	1	1	1
30 Zinc oxide 1	1	-	-	1	-
Zinc oxide 2	-	1	-	-	-
Zinc oxide 3	-	-	1	-	1
35 Palatinit	10	10	10	-	-
Total point	15	17	17	0	2
40 Zinc oxide 1: The average particle diameter is 0.5 μm and the specific surface area is 8m ² /g. Zinc oxide 2: The average particle diameter is 0.28 μm and the specific surface area is 10m ² /g. Zinc oxide 3: The average particle diameter is 0.04 μm and the specific surface area is 25m ² /g.					

* The concentration in terms of fluoride ion is 905ppm.

- 45 [0045] From the above Table 5, it was found that the astringency can be improved by decreasing the average particle diameter of zinc oxide or increasing the specific surface area thereof, and can be further improved by combining with palatinit.

Experimental Example 5

50 Organoleptic evaluation of the astringency 2

[0046] The oral compositions of Examples 17-18 and Comparative Example 10 having the varied pH values were prepared, and evaluated according to a similar manner to that described above.

[0047] The results are shown in Table 6.

Table 6

Ingr dients	Example 17	Example 18	Comparative Example 10
Silicic acid anhydride	20	20	20
Sodium carboxymethyl cellulose	1.5	1.5	1.5
Sorbitol	35	35	35
Sodium lauryl sulfate	1.5	1.5	1.5
Saccharin sodium	0.1	0.1	0.1
Sodium fluoride	0.2	0.2	0.2
Flavor	1	1	1
Zinc oxide 2*	1	1	1
Palatinit	10	10	10
Disodium hydrogenphosphate	0.1	0.1	0.1
Sodium dihydrogenphosphate	0.25	0.2	0.3
pH	6.0	7.0	5.0
Total point	16	17	10

* The average particle diameter is 0.28 μm and the specific surface area is 10m²/g.

[0048] From the above Table 6, it was shown that the astringency can be improved by adjusting pH of the oral composition to 6.0 or higher.

Example 19 Toothpaste

A toothpaste was prepared by the following formulation according to the conventional procedures.

[0049]

Ingredients	Amounts %
Palatinit	10.0
Sodium fluoride (The concentration in terms of fluoride ion is 905ppm)	0.2
Silicic acid anhydride	16.0
Sodium carboxymethyl cellulose	1.3
Sodium lauryl sulfate	1.0
Titanium dioxide	0.4
Paraoxybenzoic acid ester	0.1
Citric acid	0.1
Trisodium citrate	0.3
Saccharin sodium	0.1
Flavor	0.6
Sorbitol	50.0

EP 1 072 253 A1

(continued)

Ingredients	Amounts %
Purified water	balance
Total	100.0

Example 20 Toothpaste

[0050] A toothpaste was prepared by the following formulation according to the conventional procedures.

Ingredients	Amounts %
Palatinit	30.0
Silicic acid anhydride	20.0
Sodium carboxymethyl cellulose	1.2
Sodium lauryl sulfate	1.2
Titanium dioxide	0.3
Hydrochloric acid	0.5
Saccharin sodium	0.13
Sorbitol	10.0
Flavor	1.0
Purified water	balance
Total	100.0

Example 21 Toothpaste

[0051] A toothpaste was prepared by the following formulation according to the conventional procedures.

Ingredients	Amounts %
Palatinit	50.0
Sodium fluoride (The concentration in terms of fluoride ion is 905ppm)	0.2
Silicic acid anhydride	16.0
Carrageenan	1.3
Sodium lauryl sulfate	3.5
Titanium dioxide	0.4
Paraben	0.1
Xylitol	10.0
Flavor	0.7
Glycerol	5.0
Purified water	balance
Total	100.0

Example 22 Toothpaste

[0052] A toothpaste was prepared by the following formulation according to the conventional procedures.

Ingredients	Amounts %
Palatinit	15.0
Sodium monofluorophosphate (The concentration in terms of fluoride ion is 950ppm)	0.76
Calcium carbonate	16.0
Sodium carboxymethyl cellulose	1.3
Sodium lauroylsarcosinate	2.0
Polyoxyethylene hydrogenated castor oil	1.0
Titanium dioxide	0.4
Paraoxybenzoic acid ester	0.1
Malic acid	0.2
Stevia extract	0.1
Flavor	0.7
Sorbitol	40.0
Polyethylene glycol	5.0
Purified water	balance
Total	100.0

Example 23 Toothpaste

[0053] A toothpaste was prepared by the following formulation according to the conventional procedures.

Ingredients	Amounts %
Palatinit	5.0
Sodium fluoride (The concentration in terms of fluoride ion is 950ppm)	0.21
Triclosan	0.5
Silicic acid anhydride	16.0
Sodium polyacrylate	2.0
Sodium lauryl sulfate	1.0
Pluronic	1.0
Titanium dioxide	0.4
Paraoxybenzoic acid ester	0.1
Xylitol	10.0
Flavor	0.7
Sorbitol	50.0
Purified water	balance

(continued)

Ingredients	Amounts %
Total	100.0

Example 24 Mouthwash

[0054] A mouthwash was prepared by the following formulation according to the conventional procedures.

Ingredients	Amounts %
Palatinit	10.0
Sodium fluoride	0.05
(The concentration in terms of fluoride ion is 225ppm) Sodium lauryl sulfate	0.5
Polyoxyethylene hydrogenated castor oil	1.0
Sodium dihydrogenphosphate	0.1
Disodium hydrogenphosphate	0.1
Saccharin sodium	0.1
Flavor	0.7
Ethanol	10.0
Glycerol	10.0
Purified water	balance
Total	100.0

Example 25 Mouthwash

[0055] A mouthwash was prepared by the following formulation according to the conventional procedures.

Ingredients	Amounts %
Palatinit	30.0
Polyoxyethylene hydrogenated castor oil	1.0
Sodium citrate	0.2
Citric acid anhydride	0.2
Flavor	0.8
Glycerol	10.0
Purified water	balance
Total	100.0

Example 26 Mouthwash

[0056] A mouthwash was prepared by the following formulation according to the conventional procedures.

Ingredients	Amounts %
Palatinit	10.0
Sodium fluoride (The concentration in terms of fluoride ion is 225ppm)	0.05
Sodium lauryl sulfate	0.2
Polyoxyethylene (2)-synthetic C ₁₂ , C ₁₄ alkyl-sodium sulfosuccinate	0.2
Malic acid	0.3
Flavor	0.7
Glycerol	10.0
Xylitol	5.0
Purified water	balance
Total	100.0

Example 27 Gel dentifrice

[0057] A gel dentifrice was prepared by the following formulation according to the conventional procedures.

Ingredients	Amounts %
Palatinit	10.0
Hydroxyethyl cellulose	3.0
Sodium fluoride (The concentration in terms of fluoride ion is 4,500ppm)	1.0
Phosphoric acid	3.0
Saccharin sodium	0.5
Flavor	0.8
Glycerol	20.0
Purified water	balance
Total	100.0

Example 28 Non aerosol-type foam dentifrice

[0058] A non aerosol-type foam dentifrice was prepared by the following formulation according to the conventional procedures.

Ingredients	Amounts %
Palatinit	5.0
Sodium fluoride (The concentration in terms of fluoride ion is 4,500ppm)	1.0
Phosphoric acid	3.0
Tripotassium phosphate trihydrate	1.5
Sodium lauryl sulfate	1.0

EP 1 072 253 A1

(continued)

Ingredients	Amounts %
Pluronic	7.0
Coconut oil fatty acid diethanolamide	0.5
Saccharin sodium	0.8
Flavor	0.7
Ethanol	5.0
Purified water	balance
Total	100.0

Example 29 Oral gel

[0059] An oral gel was prepared by the following formulation according to the conventional procedures.

Ingredients	Amounts %
Palatinit	20.0
Carboxymethyl cellulose	0.2
Glycerol	40.0
Grape seed extract	1.0
α -tocopherol	0.05
Purified water	balance
Total	100.0

Example 30 Toothpaste

[0060] A toothpaste was prepared by the following formulation according to the conventional procedures.

Ingredients	Amounts %
Palatinit	10.0
Sodium fluoride (The concentration in terms of fluoride ion is 950ppm)	0.21
Silicic acid anhydride	21.0
Sodium carboxymethyl cellulose	1.1
Sodium lauryl sulfate	0.5
Sodium lauroylsarcosinate	0.1
Titanium dioxide	0.3
Fine particle zinc oxide (The average particle diameter is 0.3 μ m: the specific surface area is 10m ² /g)	1.0
Paraoxybenzoic acid ester	0.1
Saccharin sodium	0.1
Flavor	0.7
Xylitol	1.0

EP 1 072 253 A1

(continu d)

Ingredients	Amounts %
Sorbitol	38.0
Hydrochloric acid (2N)	1.0
Purified water	balance
Total	100.0
	pH 6.5

Example 31 Toothpaste

[0061] A toothpaste was prepared by the following formulation according to the conventional procedures.

Ingredients	Amounts %
Palatinit	50.0
Zinc stearate	1.0
Sodium fluoride (The concentration in terms of fluoride ion is 950ppm)	0.21
Silicic acid anhydride	16.0
Carrageenan	1.3
Sodium lauryl sulfate	3.5
Titanium dioxide	0.4
Paraben	0.1
Xylitol	10.0
Flavor	0.7
Glycerol	5.0
Purified water	balance
Total	100.0
	pH 7.0

Example 32 Toothpaste

[0062] A toothpaste was prepared by the following formulation according to the conventional procedures.

Ingredients	Amounts %
Palatinit	15.0
Zinc citrate	0.5
Sodium monofluorophosphate (The concentration in terms of fluoride ion is 950ppm)	0.75
Calcium carbonate	16.0
Sodium carboxymethyl cellulose	1.3
Sodium lauroylsarcosinate	2.0
Polyoxyethylene hydrogenated castor oil	1.0

EP 1 072 253 A1

(continued)

	Ingredients	Amounts %
5	Titanium dioxide	0.4
	Paraoxybenzoic acid ester	0.1
	Malic acid	0.2
	Stevia extract	0.1
10	Flavor	0.7
	Sorbitol	40.0
	Polyethylene glycol	5.0
	Purified water	balance
15	Total	100.0
		pH 7.5

Example 33 Toothpaste

20

[0063] A toothpaste was prepared by the following formulation according to the conventional procedures.

25	Ingredients	Amounts %
	Palatinit	5.0
	Fine particle zinc oxide (The average particle diameter is 0.3 μm : the specific surface area is 10m ² /g.)	1.5
30	Sodium fluoride (The concentration in terms of fluoride ion is 950ppm)	0.21
	Triclosan	0.5
	Silicic acid anhydride	16.0
35	Sodium polyacrylate	2.0
	Sodium lauryl sulfate	1.0
	Pluronic	1.0
	Titanium dioxide	0.4
40	Paraoxybenzoic acid ester	0.1
	Xylitol	10.0
	Flavor	0.7
45	Sorbitol	50.0
	Hydrochloric acid (1N)	1.5
	Purified water	balance
50	Total	100.0
		pH 6.8

Example 34 Mouthwash

55

[0064] A mouthwash was prepared by the following formulation according to the conventional procedures.

EP 1 072 253 A1

Ingredients	Amounts %
Palatinit	10.0
Zinc citrate	0.1
Sodium fluoride (The concentration in terms of fluoride ion is 225ppm)	0.05
Sodium lauryl sulfate	0.5
Polyoxyethylene hydrogenated castor oil	1.0
Sodium dihydrogenphosphate	0.1
Disodium hydrogenphosphate	0.1
Saccharin sodium	0.1
Flavor	0.7
Ethanol	10.0
Glycerol	10.0
Purified water	balance
Total	100.0
	pH 6.0

Example 35 Mouthwash

[0065] A mouthwash was prepared by the following formulation according to the conventional procedures.

Ingredients	Amounts %
Palatinit	10.0
Fin particle zinc oxide (The average particle diameter is 0.3 μm ; the specific surface area is 10m ² /g)	0.1
Sodium fluoride (The concentration in terms of fluoride ion is 225ppm)	0.05
Sodium lauryl sulfate	0.2
Polyoxyethylene (2)-synthetic C ₁₂ , C ₁₄ alkyl-sodium sulfosuccinate	0.2
Malic acid	0.3
Flavor	0.7
Glycerol	10.0
Xylitol	5.0
Purified water	balance
Total	100.0
	pH 7.5

Example 36 Gel dentifrice

[0066] A gel dentifrice was prepared by the following formulation according to the conventional procedures.

Ingredients	Amounts %
Palatinit	10.0
Fine particle zinc oxide (The average particle diameter is 0.3 μm : the specific surface area is 10m ² /g)	0.1
Sodium fluoride (The concentration in terms of fluoride ion is 4,500ppm)	1.0
Phosphoric acid	3.0
Saccharin sodium	0.5
Flavor	0.8
Glycerol	20.0
Hydrochloric acid (2N)	1.0
Purified water	balance
Total	100.0
	pH 6.9

Example 37 Oral gel

[0067]

Ingredients	Amounts %
Palatinit	20.0
Carboxymethyl cellulose	0.2
Fine particle zinc oxide (The average particle diameter is 0.3 μm : the specific surface area is 10m ² /g)	0.1
Glycerol	40.0
Grape seed extract	1.0
α -tocopherol	0.05
Purified water	balance
Total	100.0
	pH 8.5

Industrial Applicability

[0068] The present invention relates to an oral composition, which has the high safety, comprising either palatinit alone or in combination with fluorine compound, as well as an oral composition comprising palatinit in combination with zinc compound, which improves a bad feeling for use due to the zinc compound, and, by the inclusion of said ingredient, the present invention can provide an oral composition which can enhance remineralization.

Claims

1. An oral composition comprising palatinit.
2. The oral composition according to claim 1, further comprising a remineralization enhancing ingredient.
3. The oral composition of claim 2, comprising a fluorine compound as a remineralization enhancing ingredient.

EP 1 072 253 A1

4. The oral composition according to claim 3, wherein said fluorine compound is sodium fluoride.
5. The oral composition according to claim 2, comprising a zinc compound as a remineralization enhancing ingredient.
- 5 6. The oral composition according to claim 5, wherein said zinc compound is a water-slightly soluble zinc compound.
7. The oral composition according to claim 6, wherein said water-slightly soluble zinc compound is one or more selected from the group consisting of zinc oxide, zinc citrate and zinc stearate.
- 10 8. The oral composition according to claim 6 or 7, wherein said water-slightly soluble zinc compound is a powder having the average particle diameter of not greater than $0.3\ \mu\text{m}$ and the specific surface area of greater than $10\text{m}^2/\text{g}$.
9. The oral composition according to any one of claims 5-8, which as pH of 6.0-8.5.
- 15 10. The oral composition according to claim 9, wherein hydrochloric acid is used as a pH adjusting agent.
11. The oral composition according to any one of claims 5-10, further comprising a fluorine compound.
- 20 12. The oral composition of claim 11, wherein said fluorine compound is sodium fluoride.

Figure 1





Figure 2



INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP99/02060

A. CLASSIFICATION OF SUBJECT MATTER Int.Cl. ⁶ A61K7/16		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int.Cl. ⁶ A61K7/16		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAPLUS (STN)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category ^a	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JP, 6-70704, A (Towa Chemical Industry Co., Ltd.), 15 March, 1994 (15. 03. 94), Claims (Family: none)	1
X	JP, 7-132047, A (Lotte Co., Ltd.), 23 May, 1995 (23. 05. 95), Claims & US, 5629040, A	1
X	JP, 8-214774, A (Yakult Honsha Co., Ltd.), 27 August, 1996 (27. 08. 96), Claims & WO, 9625050, A1 & EP, 809939, A1	1
X	JP, 9-199, A (Meiji Seika Kaisha, Ltd.), 7 January, 1997 (07. 01. 97), Claims ; Par. No. [0013] (Family: none)	1
X	JP, 9-107900, A (Towa Chemical Industry Co., Ltd.), 28 April, 1997 (28. 04. 97), Claims ; Par. Nos. [0037], [0038] (Family: none)	1
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
^a Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 12 July, 1999 (12. 07. 99)		Date of mailing of the international search report 21 July, 1999 (21. 07. 99)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP99/02060

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category ^a	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JP, 9-238642, A (Towa Chemical Industry Co., Ltd.), 16 September, 1997 (16. 09. 97), Claims ; Par. No. [0004] (Family: none)	1
PX	JP, 10-155410, A (Snow Brand Milk Products Co., Ltd.), 16 June, 1998 (16. 06. 98), Claims ; Par. No. [0008] (Family: none)	1
A	JP, 60-204710, A (Lion Corp.), 16 October, 1985 (16. 10. 85), Full text (Family: none)	1-12

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

Figure 1




Figure 2

